

The end of the Ediacaran: Two new exceptionally preserved body fossil assemblages from Mount Dunfee, Nevada, USA

E.F. Smith¹, L.L. Nelson², M.A. Strange³, A.E. Eyster², S.M. Rowland³, D.P. Schrag², and F.A. Macdonald²

¹Department of Paleobiology, Smithsonian Institution, PO Box 37012, MRC 121, Washington, D.C. 20013-7012, USA

²Department of Earth and Planetary Sciences, Harvard University, 20 Oxford Street, Cambridge, Massachusetts 02138, USA

³Department of Geoscience, University of Nevada, Las Vegas, 4505 S. Maryland Parkway, Las Vegas, Nevada 89154-4010, USA

ABSTRACT

Evaluation of hypotheses that relate environmental to evolutionary change across the Ediacaran-Cambrian transition has been hampered by a dearth of sections that preserve both the last appearance of Ediacaran body fossils and the first appearance of *Treptichnus pedum* within carbonate-rich strata suitable for chemostratigraphic studies. Here, we report two new exceptionally preserved latest Ediacaran fossil assemblages from the Deep Spring Formation at Mount Dunfee, Nevada (USA). Further, we report these occurrences in a high-resolution carbon isotope chemostratigraphic framework, permitting correlation on a regional and global scale. The lower of the two horizons, at the base of the Deep Spring Formation, hosts a body fossil assemblage that includes *Gaojiashania*, other vermiform body fossils, and possible *Wutubus annularis* interbedded with *Cloudina* shell beds. The upper of the two fossil horizons, in the Esmeralda Member of the Deep Spring Formation, contains *Conotubus* and occurs within the basal Cambrian negative carbon isotope excursion, establishing it as the youngest Ediacaran fossil assemblage discovered to date. This is the first report of *Gaojiashania*, *Conotubus*, and *Wutubus* in Laurentia, extending the known stratigraphic ranges and biogeographic distributions of these taxa to a global scale. These data refine the relative ages of defining characteristics of the Ediacaran-Cambrian boundary and confirm that a large perturbation to the carbon cycle and surface ocean conditions coincided with the extinction of Ediacaran organisms.

INTRODUCTION

The first major radiation of macroscopic multicellular life occurred during the Ediacaran Period. Although Ediacaran organisms likely represent a diverse assemblage of eukaryotic groups—including stem- and crown-group metazoans, algae, and extinct clades with no modern representatives—they are commonly collectively referred to as the “Ediacara biota” (Xiao and Laflamme, 2009; Erwin et al., 2011; Brasier et al., 2012). Across the Ediacaran-Cambrian transition, both macroscopic soft-bodied Ediacara biota and enigmatic, tubular metazoan biota (Cai et al., 2013; Tarhan et al., 2014) disappear from the fossil record. Although it has been suggested that the apparent extinction of Ediacaran organisms is a taphonomic artifact (Gehling, 1991), the pattern of biological turnover appears to be globally reproducible through a wide range of sedimentary facies (Buatois et al., 2014). There are a few isolated reports of purported “Cambrian survivors” of Ediacaran taxa (Hagadorn et al., 2000; Jensen et al., 1998), but more taphonomic and taxonomic studies are necessary to demonstrate that they are in fact true Ediacaran holdovers (Laflamme et al., 2013).

Two general classes of hypotheses have been proposed to explain the disappearance of the Ediacaran organisms: (1) an extreme perturbation to

surface environments and global geochemical cycles marked by a carbon isotope ($\delta^{13}\text{C}$) excursion triggered an extinction by generating abiotic stressors (e.g., Knoll and Carroll, 1999; Amthor et al., 2003), and (2) a growing metazoan community more gradually outcompeted Ediacaran soft-bodied non-metazoan taxa through ecological engineering and gradual biotic replacement (e.g., Erwin and Tweedt, 2012; Darroch et al., 2015). Calibrating the relative and absolute timing of changes in life and environment at this interval—the necessary framework to test these hypotheses—has been hindered by a lack of fossiliferous, carbonate-rich sections in which both paleontological and geochemical proxies can be integrated. Here we present new geological mapping, stratigraphic sections, geochemical data, and two newly discovered assemblages of Ediacaran body fossils from Mount Dunfee, Nevada (USA). These data allow us to directly relate the nadir of the multi-pronged basal Cambrian negative $\delta^{13}\text{C}$ excursion to the last appearance datum (LAD) of Ediacaran body fossils and the first appearance datum (FAD) of *Treptichnus pedum*, which defines the base of the Cambrian Period.

LITHOSTRATIGRAPHY

Ediacaran-Cambrian strata in the Great Basin of Nevada and California were deposited on the

western margin of Laurentia (present coordinates) during the rift-drift transition (Armin and Mayer, 1983; Stewart, 1970), with the strata at Mount Dunfee (Figs. 1A and 1B) representing one of the most distal, carbonate-dominated sections that is preserved. The Ediacaran to lower Cambrian Deep Spring Formation consists of three members: the Dunfee (lower), Esmeralda (middle), and Gold Point (upper) Members, which are also exposed in the White-Inyo Mountains (Ahn et al., 2012; Nelson, 1962).

At Mount Dunfee, the contact between the Reed Dolomite and the overlying Deep Spring Formation has previously been placed at the transition from thickly bedded dolostone of the Reed Dolomite to distinctly bedded dolostone, limestone, and fine-grained siliciclastics of the Deep Spring Formation (Albers and Stewart, 1972). Here, because this gradational contact coincides with an irregular dolomitization front and lateral facies change, we refer to the bedded orange and gray dolostone and limestone beds above the massive, structureless dolostone as the “Reed–Deep Spring transition beds.” At Mount Dunfee, these transition beds contain ooids, pisoids, oncoids, and shelly fossils in lenticular lag deposits (Gevirtzman and Mount, 1986).

We place the base of the Deep Spring Formation at a regionally distinct and easily recognizable sequence boundary (Fig. 1C). This sequence boundary is a sharp, karsted contact between a red, Fe-rich, glauconitic, peloidal limestone and a fine-grained siliciclastic facies (interbedded shale, siltstone, and sandstone) that contains mud chips, mud cracks, and syneresis cracks. The rest of the Dunfee Member consists of mixed carbonate-siliciclastic deposits representing slope to shallow subtidal settings (Gevirtzman and Mount, 1986).

The contact between the Dunfee and Esmeralda Members is marked by a pink to gray recrystallized dolostone capped by a karstic dissolution surface that is sharply overlain by green shoreface sandstone with mud cracks, interference ripples, and bed-planar trace fossils. The Esmeralda Member is 275–325 m thick and is composed of quartzitic and calcareous sandstone, stromatolitic and oolitic limestone, and minor siltstone and shale (Albers and Stewart, 1972).




Figure 1. A: Locality map for Mount Dunfee, Nevada, USA (LA—Los Angeles, California; LV—Las Vegas, Nevada). **B:** Geologic map of Mount Dunfee with locations of measured sections. Section E1513 is northwest of figured map area (Base: 37°21'1.42"N, 117°19'37.85"W). **C:** Composite measured section with biostratigraphy and δ¹³C chemostratigraphy for the upper Reed Dolomite and Deep Spring Formation. Stratigraphic horizons of two new Ediacaran body fossil assemblages are marked with stars. DSF—Deep Spring Formation; VPDB—Vienna Pee Dee belemnite. vf-f ss—very fine to fine-grained sandstone; m ss—medium-grained sandstone; c-vc ss—coarse to very coarse-grained sandstone; lm/ds—limestone and dolostone; cong—conglomerate.

CHEMOSTRATIGRAPHY

Detailed geologic mapping of Miocene down-to-the-southwest shingled normal faults in the Mount Dunfee area (Fig. 1B) and correlation of distinct marker beds enabled the construction of a composite measured section and a high-resolution δ¹³C chemostratigraphic curve (see the GSA Data Repository¹) of the upper Reed Dolomite through the Esmeralda Member

¹GSA Data Repository item 2016307, methods, measured sections, and carbon and oxygen data tables, is available online at www.geosociety.org/pubs/ft2016.htm, or on request from editing@geosociety.org.

of the Deep Spring Formation (Fig. 1C). At Mount Dunfee, the upper Reed Dolomite and lower Deep Spring Formation are characterized by a +3‰ to +4‰ δ¹³C plateau (Fig. 1C). Above this interval, within the dolomitized and recrystallized upper ~50 m of the Dunfee Member, there is a downturn in δ¹³C values to -2‰. The previously documented basal Cambrian negative δ¹³C excursion (Corsetti and Kaufman, 1994) was reproduced in the middle of the Esmeralda Member, but new high-resolution data reveal additional structure to the excursion (Fig. 1C). Conglomerate with tabular carbonate intraclasts and microbial horizons A and B (Oliver and

Rowland, 2002) contain δ¹³C values as low as -6.2‰. Above this, the δ¹³C values rise to -3‰ to -2‰ before they decrease to -9.5‰.

BIOSTRATIGRAPHY

Small shelly fossils were previously reported from the upper Reed Dolomite and the lower Dunfee Member of the Deep Spring Formation (Gevirtzman and Mount, 1986; Signor et al., 1987) and were later reinterpreted as *Cloudina* (Grant, 1990). In this study, shelly debris with circular cross-sections characteristic of *Cloudina* were discovered in the uppermost bed of the Dunfee Member, extending, in a chemostratigraphic and biostratigraphic context, the LAD of this Ediacaran index fossil (Fig. 1C). The algal fossil *Elainabella deepspringensis* occurs in brown to black siltstone above microbial horizon B and below microbial horizon C of the Esmeralda Member (Rowland and Rodriguez, 2014). The FAD of *T. pedum* occurs above the large negative δ¹³C excursion, just a few meters below the top of the Esmeralda Member (Corsetti and Hagadorn, 2003). Additionally, two new exceptionally preserved Ediacaran body fossil assemblages were discovered in the Deep Spring Formation and are described in detail below.

Fossil Assemblage in the Dunfee Member of the Deep Spring Formation

The older of the two body fossil assemblages reported here occurs in siltstone and sandstone beds in the lowermost Dunfee Member of the Deep Spring Formation (Fig. 1C). Cast and mold impressions of annulated tubular fossils, identified as *Gaojiashania*, were discovered on multiple bedding surfaces within a 5–10-m-thick interval of strata composed of interbedded green shale, micaceous brown to green siltstone, and micaceous fine- to medium-grained sandstone (Figs. 2A and 2B). The specimens are 1.5–10 cm in length and 3–13 mm in diameter. Other fossils in this stratigraphic interval include cast and mold impressions of possible *Wutubus annularis* (Fig. 2C), lightly pyritized compressions of smaller, tubular vermicular fossils with closed apices (Fig. 2D), and possible algal fossils. Part and counterpart slabs were collected for five *Gaojiashania* specimens. Some of the fossiliferous horizons contain <1-cm-diameter subangular to rounded mud chips. Mud cracks, syneresis cracks, and hummocky cross-stratification are present both within and above the fossiliferous interval. Small (submillimeter- to millimeter-scale) trace fossils, including *Planolites*, *Bergaueria*-like traces, and rare traces resembling *Scolicia* and *Neonereites*, occur in this same interval (Gevirtzman and Mount, 1986). Body fossils described herein were recovered from three separate fault blocks around Mount Dunfee.

To date, all reports of *Gaojiashania* are from the late Ediacaran (ca. 551–541 Ma) Dengying




Figure 2. Photographs of fossils from the lower Dunfee Member of the Deep Spring Formation, Nevada, USA. A: Cast of *Gaojiashania* specimen preserved in siltstone. B: *Gaojiashania* specimen that preserves inner and outer fossil walls. C: Elongate, tapering, annulated fossils that we suggest should be assigned to the genus *Wutubus*. D: One of four vermicular-like body fossil specimens.

Formation (Cai et al., 2010; Lin et al., 1986) and Taozhong Formation (Hua et al., 2004) of South China. Like the specimens from South China, the Mount Dunfee specimens occur as compressed cylinders with densely spaced, transverse rings that appear to be articulated by a very thin membrane (Figs. 2A and 2B). Cai et al. (2011a) pointed to this feature as a way of distinguishing *Gaojiashania* from the late Ediacaran ribbon-shaped fossil *Shaanxilithes ningqiangensis* that is composed of serially arranged discoidal units (Meyer et al., 2012). However, more recently, Tarhan et al. (2014) interpreted the transverse ridges of *Shaanxilithes* specimens from India as representing anatomical wall thickenings of a tubular organism. On the basis of morphological characteristics, we classify the Mount Dunfee fossils as *Gaojiashania*, but acknowledge that these two fossil genera could be the same organism.

Three specimens of an additional annulated tubular taxon, which differ in size and morphology from *Gaojiashania*, also occur in the Dunfee Member assemblage. These specimens are characterized by a narrower conical termination and are similar in size and morphology to *Wutubus annularis* specimens from the Shibantan Member (correlative with the Gaojiashan Member) of the Dengying Formation of South China (Chen et al., 2014) (Fig. 2C).

Fossil Assemblage in the Esmeralda Member of the Deep Spring Formation

The upper of the two fossiliferous horizons (Fig. 1C) at Mount Dunfee occurs in the middle part of the Esmeralda Member of the Deep

Spring Formation in shale and siltstone beds between microbial horizons B and C (Oliver and Rowland, 2002). These fossils consist of narrow conical tubes with transverse annulations (Figs. 3A and 3B). The diameter of these tubular fossils is ~1 mm and the length of articulated specimens ranges from 0.5 to 3 cm. Cross-sections of these fossils are circular, and the narrow end of these fossils is tapered and closed (Fig. 3A). Non-uniform curvature is present in some of these fossils, and many of the specimens are characterized by funnel-in-funnel structure that is characteristic of the family Cloudinidae (Figs. 3A and 3B). These fossils are commonly stained orange or red, in contrast to the black, brown, or green matrix of the host siltstone and




Figure 3. Photographs of pyritized *Conotubus* specimen from the Esmeralda Member of the Deep Spring Formation, Nevada, USA. A: *Conotubus* specimen that exhibits bending and high-relief preservation. B: High-resolution photograph of the same specimen that shows funnel-in-funnel structure.

shale, most likely reflecting oxidation of a pyrite pseudomorph remnant of the original fossil wall. This same stratigraphic interval also contains the multicellular algae *Elainabella deepspringensis* (Rowland and Rodriguez, 2014).

Based on morphological similarity, we classify these annulated tubular fossils as *Conotubus*, which, like *Gaojiashania*, until now has been described only from the Gaojiashan Lagerstätte in South China (Cai et al., 2011a; Hua et al., 2007). The pyritization of the Dunfee specimens is also similar to that of the *Conotubus* fossils in South China (Cai et al., 2012; Schiffbauer et al., 2014). Like *Cloudina*, *Conotubus* is characterized by tubular morphology with nested cylindrical or funnel-shaped units, but what distinguishes the two genera is that *Cloudina* are interpreted to have been biomineralized (Grant, 1990), whereas *Conotubus* are either weakly biomineralized or non-biomineralized (Hua et al., 2007). The combination of non-uniform bends and the high relief of some of the fossil specimens described herein (Fig. 3A) suggests that the original wall could have been lightly calcified or composed of a resistant biomolecule such as chitin. Due to similarities in morphology and interpreted epibenthic life mode, *Conotubus* has been interpreted as a potential evolutionary precursor to *Cloudina* (Cai et al., 2011b; Hua et al., 2007). The data presented here, however, establish an inverse biostratigraphic relationship between *Conotubus* and *Cloudina*, in which *Conotubus* appears ~100 m above the LAD of *Cloudina* (Fig. 1C).

DISCUSSION

Two new, exceptionally preserved late Ediacaran body fossil assemblages at Mount Dunfee compose the only known Ediacaran Lagerstätten in the southwestern United States. Discovery of these fossil assemblages expands the biogeographic distributions and stratigraphic ranges of the tubular fossils *Gaojiashania*, *Conotubus*, and *Wutubus*, adding another late Ediacaran fossil locality to the global record, and specifically, one dominated by tubular metazoans. Furthermore, at Mount Dunfee, the biostratigraphic range of *Gaojiashania* overlaps with that of *Cloudina*, instead of occurring exclusively below *Cloudina* as it does in South China (Cai et al., 2010).

The two competing models for the cause of the end-Ediacaran extinction—an exogenous shock to surface environments or more gradual biotic replacement of soft-bodied Ediacaran organisms (e.g., erniettomorphs, rangeomorphs) by tubular metazoans—are not mutually exclusive, and earlier stages of biotic replacement could have preceded an ultimate extinction event. However, the data set presented here shows that the nadir of the $\delta^{13}\text{C}$ excursion, the LAD of Ediacaran tubular fossils, and the FAD of *T. pedum*, which defines the Precambrian-Cambrian

boundary, all occur within ~100 m of stratigraphy. The large $\delta^{13}\text{C}$ excursion likely records a major perturbation to the carbon cycle through the addition of isotopically light carbon to the ocean-atmosphere system, and possibly concomitant acidification of the surface ocean. The data presented here represent the tightest relationship documented to date between the negative $\delta^{13}\text{C}$ excursion and biological turnover at the Ediacaran-Cambrian boundary, consistent with an environmental disturbance eliminating the last of the Ediacaran biota and paving the way for the Cambrian radiation.

ACKNOWLEDGMENTS

We thank the National Science Foundation Graduate Research Fellowship Program and the Massachusetts Institute of Technology NASA Astrobiology Institute for support; E. Hodgkin and F. Liu for assistance in the field; E. Sperling, L. Tarhan, D. Erwin, and S. Tweedt for advice and stimulating discussions; and J. Schiffbauer, M. Laflamme, and two anonymous reviewers for improving this manuscript.

REFERENCES CITED

- Ahn, S.Y., Babcock, L.E., and Hollingsworth, J., 2012, Revised stratigraphic nomenclature for parts of the Ediacaran-Cambrian Series 2 succession in the southern Great Basin, USA: *Memoirs of the Association of Australasian Palaeontologists*, v. 42, p. 105–114.
- Albers, J.P., and Stewart, J.H., 1972, Geology and mineral deposits of Esmeralda County, Nevada: Nevada Bureau of Mines and Geology Bulletin 78, 88 p., 2 plates, scale 1:250,000.
- Amthor, J.E., Grotzinger, J.P., Schröder, S., Bowring, S.A., Ramezani, J., Martin, M.W., and Matter, A., 2003, Extinction of *Cloudina* and *Namacalathus* at the Precambrian-Cambrian boundary in Oman: *Geology*, v. 31, p. 431–434, doi:10.1130/0091-7613(2003)031<0431:EOCANA>2.0.CO;2.
- Armin, R.A., and Mayer, L., 1983, Subsidence analysis of the Cordilleran miogeocline: Implications for timing of late Proterozoic rifting and amount of extension: *Geology*, v. 11, p. 702–705, doi:10.1130/0091-7613(1983)11<702:SAOTCM>2.0.CO;2.
- Brasier, M.D., Antcliffe, J.B., and Liu, A.G., 2012, The architecture of Ediacaran fronds: *Palaeontology*, v. 55, p. 1105–1124, doi:10.1111/j.1475-4983.2012.01164.x.
- Buatois, L.A., Narbonne, G.M., Mángano, M.G., Carmona, N.B., and Myrow, P., 2014, Ediacaran mat-ground ecology persisted into the earliest Cambrian: *Nature Communications*, v. 5, 3544, doi:10.1038/ncomms4544.
- Cai, Y., Hua, H., Xiao, S., Schiffbauer, J.D., and Li, P., 2010, Biostratigraphy of the late Ediacaran pyritized Gaojiashan Lagerstätte from southern Shaanxi, South China: Importance of event deposits: *Palaios*, v. 25, p. 487–506, doi:10.2110/palo.2009.p09-133r.
- Cai, Y., Hua, H., Zhuravlev, A.Y., Gámez Vintaned, J.A., and Ivantsov, A.Y., 2011a, Discussion of 'First finds of problematic Ediacaran fossil *Gaojiashania* in Siberia and its origin': *Geological Magazine*, v. 148, p. 329–333, doi:10.1017/S0016756810000749.
- Cai, Y., Schiffbauer, J.D., Hua, H., and Xiao, S., 2011b, Morphology and paleoecology of the late Ediacaran tubular fossil *Conotubus hemiannulatus* from the Gaojiashan Lagerstätte of southern Shaanxi Province, South China: *Precambrian Research*, v. 191, p. 46–57, doi:10.1016/j.precamres.2011.09.002.
- Cai, Y., Schiffbauer, J.D., Hua, H., and Xiao, S., 2012, Preservation modes in the Ediacaran Gaojiashan Lagerstätte: Pyritization, aluminosilicification, and carbonaceous compression: *Palaeogeography, Palaeoclimatology, Palaeoecology*, v. 326, p. 109–117, doi:10.1016/j.palaeo.2012.02.009.
- Cai, Y., Hua, H., and Zhang, X., 2013, Tube construction and life mode of the late Ediacaran tubular fossil *Gaojiashania cyclus* from the Gaojiashan Lagerstätte: *Precambrian Research*, v. 224, p. 255–267, doi:10.1016/j.precamres.2012.09.022.
- Chen, Z., Zhou, C., Xiao, S., Wang, W., Guan, C., Hua, H., and Yuan, X., 2014, New Ediacara fossils preserved in marine limestone and their ecological implications: *Scientific Reports*, v. 4, 4180, doi:10.1038/srep04180.
- Corsetti, F.A., and Hagadorn, J.W., 2003, The Precambrian-Cambrian transition in the southern Great Basin, USA: *The Sedimentary Record*, v. 1, p. 4–8.
- Corsetti, F.A., and Kaufman, A.J., 1994, Chemostratigraphy of Neoproterozoic-Cambrian units, White-Inyo Region, eastern California and western Nevada: Implications for global correlation and faunal distribution: *Palaios*, v. 9, p. 211–219, doi:10.2307/3515107.
- Darroch, S.A.F., Sperling, E.A., Boag, T.H., Racicot, R.A., Mason, S.J., Morgan, A.S., Tweedt, S., Myrow, P., Johnston, D.T., and Erwin, D.H., 2015, Biotic replacement and mass extinction of the Ediacara biota: *Proceedings of the Royal Society of London B*, v. 282, 20151003, doi:10.1098/rspb.2015.1003.
- Erwin, D.H., and Tweedt, S., 2012, Ecological drivers of the Ediacaran-Cambrian diversification of Metazoa: *Evolutionary Ecology*, v. 26, p. 417–433, doi:10.1007/s10682-011-9505-7.
- Erwin, D.H., Laflamme, M., Tweedt, S.M., Sperling, E.A., Pisani, D., and Peterson, K.J., 2011, The Cambrian conundrum: Early divergence and later ecological success in the early history of animals: *Science*, v. 334, p. 1091–1097, doi:10.1126/science.1206375.
- Gehling, J.G., 1991, The case for Ediacaran fossil roots to the metazoan tree: *Geological Society of India Memoir*, v. 20, p. 181–224.
- Gevirtzman, D.A., and Mount, J.F., 1986, Paleoenvironments of an earliest Cambrian (Tommotian) shelly fauna in the southwestern Great Basin, USA: *Journal of Sedimentary Petrology*, v. 56, p. 412–421.
- Grant, S.W., 1990, Shell structure and distribution of *Cloudina*, a potential index fossil for the terminal Proterozoic: *American Journal of Science*, v. 290, p. 261–294.
- Hagadorn, J.W., Fedo, C.M., and Waggoner, B.M., 2000, Early Cambrian Ediacaran-type fossils from California: *Journal of Paleontology*, v. 74, p. 731–740, doi:10.1017/S0022336000032832.
- Hua, H., Chen, Z., and Zhang, L., 2004, *Shaanxilithes* from lower Taozichong Formation, Guizhou Province and its geological and paleobiological significance: *Journal of Stratigraphy*, v. 28, p. 265–269.
- Hua, H., Chen, Z., and Yuan, X., 2007, The advent of mineralized skeletons in Neoproterozoic Metazoa—New fossil evidence from the Gaojiashan Fauna: *Geological Journal*, v. 42, p. 263–279, doi:10.1002/gj.1077.
- Jensen, S., Gehling, J.G., and Droser, M.L., 1998, Ediacara-type fossils in Cambrian sediments: *Nature*, v. 393, p. 567–569, doi:10.1038/31215.
- Knoll, A.H., and Carroll, S.B., 1999, Early animal evolution: Emerging views from comparative biology and geology: *Science*, v. 284, p. 2129–2137, doi:10.1126/science.284.5423.2129.
- Laflamme, M., Darroch, S.A.F., Tweedt, S.M., Peterson, K.J., and Erwin, D.H., 2013, The end of the Ediacara biota: Extinction, biotic replacement, or Cheshire Cat?: *Gondwana Research*, v. 23, p. 558–573, doi:10.1016/j.gr.2012.11.004.
- Lin, S., Zhang, Y., Zhang, L., Tao, X., and Wang, M., 1986, Body and trace fossils of Metazoa and algal macrofossils from the upper Sinian Gaojiashan Formation in southern Shaanxi: *Geology of Shaanxi*, v. 4, p. 9–17.
- Meyer, M., Schiffbauer, J.D., Xiao, S., Cai, Y., and Hua, H., 2012, Taphonomy of the upper Ediacaran enigmatic ribbonlike fossil *Shaanxilithes*: *Palaios*, v. 27, p. 354–372, doi:10.2110/palo.2011.p11-098r.
- Nelson, C.A., 1962, Lower Cambrian-Precambrian succession, White-Inyo Mountains, California: *Geological Society of America Bulletin*, v. 73, p. 139–144, doi:10.1130/0016-7606(1962)73[139:LCSWMC]2.0.CO;2.
- Oliver, L.K., and Rowland, S.M., 2002, Microbialite reefs at the close of the Proterozoic Eon: The Middle Member Deep Spring Formation at Mt. Dunfee, Nevada, in Corsetti, F.A., ed., *Proterozoic-Cambrian of the Great Basin and Beyond: SEPM (Society for Sedimentary Geology) Pacific Section Book 93*, p. 97–122.
- Rowland, S.M., and Rodriguez, M.G., 2014, A multicellular alga with exceptional preservation from the Ediacaran of Nevada: *Journal of Paleontology*, v. 88, p. 263–268, doi:10.1666/13-075.
- Schiffbauer, J.D., Xiao, S., Cai, Y., Wallace, A.F., Hua, H., Hunter, J., Xu, H., Peng, Y., and Kaufman, A.J., 2014, A unifying model for Neoproterozoic–Palaeozoic exceptional fossil preservation through pyritization and carbonaceous compression: *Nature Communications*, v. 5, 5754, doi:10.1038/ncomms6754.
- Signor, P.W., Mount, J.F., and Onken, B.R., 1987, A pre-trilobite shelly fauna from the White-Inyo region of eastern California and western Nevada: *Journal of Paleontology*, v. 61, p. 425–438, doi:10.1017/S0022336000028614.
- Stewart, J.H., 1970, Upper Precambrian and Lower Cambrian strata in the southern Great Basin, California and Nevada: U.S. Geological Survey Professional Paper 620, 206 p.
- Tarhan, L.G., Hughes, N.C., Myrow, P.M., Bhargava, O.N., Ahluwalia, A.D., and Kudryavtsev, A.B., 2014, Precambrian-Cambrian boundary interval occurrence and form of the enigmatic tubular body fossil *Shaanxilithes ningqiangensis* from the Lesser Himalaya of India: *Palaeontology*, v. 57, p. 283–298, doi:10.1111/pala.12066.
- Xiao, S., and Laflamme, M., 2009, On the eve of animal radiation: Phylogeny, ecology and evolution of the Ediacara biota: *Trends in Ecology & Evolution*, v. 24, p. 31–40, doi:10.1016/j.tree.2008.07.015.

Manuscript received 27 May 2016

Revised manuscript received 12 August 2016

Manuscript accepted 15 August 2016

Printed in USA